



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

AVERAGE AND PROBABILITY.

Conducted by B. F. FINKEL, Springfield, Mo. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.


30. Proposed by F. P. MATZ, M. A., M. Sc., Ph. D., Professor of Mathematics and Astronomy in Irving College, Mechanicsburg, Pennsylvania.

Find the average area of all the triangles which can be inscribed in a given circle.

I. Solution by the PROPOSER.

Let P_1OP_2 be any inscribed triangle; and through O draw any diameter OA . Two cases have now to be considered: (1), the triangle may lie *wholly* on one side of the diameter OA ; (2), the triangle may lie *partly* on one side of the diameter OA .

I. Put $OA=2r$, $\angle AOP_1=\phi$, and $\angle AOP_2=\theta$; then $OP_1=2r\cos\phi$, $OP_2=2r\cos\theta$, and the area of the $\triangle, P_1OP_2, =A', =2r^2\cos\phi\cos\theta\sin(\phi-\theta)$. Hence the average area of the triangles in this case, is



$$A_1 = \int_0^{\frac{1}{2}\pi} \int_0^{\phi} A' d\phi d\theta \div \int_0^{\frac{1}{2}\pi} \int_0^{\phi} d\phi d\theta = \frac{8r^2}{\pi^2} \int_0^{\frac{1}{2}\pi} \phi \sin\phi \cos\phi d\phi \\ = \frac{r^2}{\pi^2} \left[\sin 2\phi - 2\phi \cos 2\phi \right]_0^{\frac{1}{2}\pi} = \frac{r^2}{\pi} \dots \dots \dots (1).$$

II. Put $\angle AOP_3=\psi$; then the area of the triangle $P_2OP_3, =A'', =2r^2\cos\theta\cos\psi\sin(\theta+\psi)$. Hence the average area of the triangles in this case, is

$$A_2 = \int_0^{\frac{1}{2}\pi} \int_0^{\frac{1}{2}\pi} A'' d\theta d\psi \div \int_0^{\frac{1}{2}\pi} \int_0^{\frac{1}{2}\pi} d\theta d\psi = \frac{8r^2}{\pi^2} \left[\frac{\pi}{4} \int_0^{\frac{1}{2}\pi} \sin\theta \cos\theta d\theta \right. \\ \left. + \frac{1}{2} \int_0^{\frac{1}{2}\pi} \cos^2\theta d\theta \right] = \frac{8r^2}{\pi^2} \left[\frac{\pi}{4} \times \frac{1}{2} + \frac{1}{2} \times \frac{\pi}{4} \right] = \frac{2r^2}{\pi} \dots \dots \dots (2).$$

Hence the *required* average area becomes

$$A = \frac{1}{2}(A_1 + A_2) = 3r^2 / 2\pi \dots \dots \dots (3).$$

II. Solution by O. W. ANTHONY, M. Sc., Professor of Mathematics and Astronomy, New Windsor College, New Windsor, Maryland.

We readily get the area of triangle

$$= \frac{R^2}{2} \{ \sin 2A + \sin 2B + \sin 2C \},$$

which, by virtue of the relation $A+B+C=\pi$, reduces to

$$\frac{R^2}{2} \{\sin 2A + \sin 2B - \sin 2(A+B)\}.$$

$$\therefore \text{Average area} = \frac{R^2}{2} \frac{\int_0^\pi \int_0^{\pi-A} \{\sin 2A + \sin 2B - \sin 2(A+B)\} dA dB}{\int_0^\pi \int_0^{\pi-A} dA dB} = \frac{3R^2}{2\pi}.$$

31. Proposed by B. F. FINKEL, A. M., Professor of Mathematics and Physics in Drury College, Springfield, Missouri.

Find the average length of a line drawn across the opposite sides of a rectangle, length l and breadth b .

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Applied Science, Texarkana College, Texarkana, Arkansas-Texas; and the PROPOSER.

Let $ABCD$ be the rectangle, FG the random line. Let $AB=l$, $BC=b$, $AH=x$, $AG=y$.

Then $FG = \{b^2 + (x-y)^2\}^{\frac{1}{2}}$.

The limits of x are 0 and l ; of y , 0 and x .

Hence the required average area is

$$A = \frac{\int_0^l \int_0^x \{b^2 + (x-y)^2\}^{\frac{1}{2}} dx dy}{\int_0^l \int_0^x dx dy}$$

$$= \frac{2}{l^2} \int_0^l \int_0^x \{b^2 + (x-y)^2\}^{\frac{1}{2}} dx dy$$

$$= \frac{1}{l^2} \int_0^l \{x(b^2 + x^2)^{\frac{1}{2}} + b^2 \log[x + (b^2 + x^2)^{\frac{1}{2}}] - b^2 \log b\} dx$$

$$= \frac{1}{3l^2} (l^2 + b^2)^{\frac{3}{2}} + \frac{b^2}{l} \log\{l + (l^2 + b^2)^{\frac{1}{2}}\} - \frac{b^2}{l} \log b - \frac{1}{l^2} (l^2 + b^2)^{\frac{1}{2}} - \frac{b^3}{3l^2} + \frac{b}{l^2}.$$

For the line KL , we get, by writing l for b and b for l ,

$$A_1 = \frac{1}{3b^2} (l^2 + b^2)^{\frac{3}{2}} + \frac{l^2}{b} \log\{b + (l^2 + b^2)^{\frac{1}{2}}\} - \frac{l^2}{b} \log l - \frac{1}{b^2} (l^2 + b^2)^{\frac{1}{2}} - \frac{l^3}{3b^2} + \frac{l}{b^2}.$$

$$\text{Cor. I. If } l=b, A = \frac{1}{3}(2l\sqrt{2}) + l \log(1+\sqrt{2}) - \frac{1}{l}\sqrt{2} - \frac{1}{3}l + \frac{1}{l}.$$

Cor. II. If $l=b=1$, $A = \frac{1}{3}(2-\sqrt{2}) + \log(1+\sqrt{2})$, which is the same result as given in *Williamson's Integral Calculus*, page 409.

Also solved by F. P. MATZ.

